

Surname	
Other Names	
Centre Number	
Candidate Number	
Candidate Signature	
I declare this is my own wor	k.

A-level BIOLOGY

Paper 3

7402/3

Time allowed: 2 hours

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.



For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Answer ALL questions in SECTION A.
- Answer ONE question from SECTION B.
- You must answer the questions in the spaces provided. Do not write on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.



INFORMATION

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 78.

DO NOT TURN OVER UNTIL TOLD TO DO SO



BLANK PAGE



SECTION A

Answer ALL questions in this section.

You are advised to spend no more than 1 hour and 15 minutes on this section.



0 1

In one species of squirrel, 'Sciurus carolinensis', fur colour is controlled by one gene, with two codominant alleles. C^G represents the allele for grey fur colour, and C^B represents the allele for black fur colour.

TABLE 1 shows the three possible phenotypes.

TABLE 1

GENOTYPE	PHENOTYPE
CGCG	Grey fur
CGCB	Brown-black fur
CBCB	Black fur



0 1.	. 1
------	-----

In a population of 34 'S. carolinensis', 2 had black fur.

Use the Hardy–Weinberg equation to estimate how many squirrels in this population had brown-black fur. Show your working. [2 marks]



0 1 . 2

The actual number of squirrels in this population that had brown-black fur was 16.

Use all of the information to calculate the ACTUAL frequency of the C^G allele.

Do NOT use the Hardy–Weinberg equation in your calculation.

Give your answer to 2 decimal places. [1 mark]

Answer ____



BLANK PAGE



0 1 . 3

- 'S. carolinensis' were first introduced to the UK from North America in the 1870s. They are now widely distributed across the UK.
- 'S. carolinensis' from both North America and the UK show exactly the same genotypic and phenotypic variation. An identical mutation causing black fur has also been found in several other species closely related to 'S. carolinensis'.

Use this information to deduce which ONE of the following conclusions is most likely true.

On the opposite page, tick (√) ONE box. [1 mark]



The mutation that caused black fur happened after 'S. carolinensis' was introduced to the UK from North America.
The mutation that caused black fur happened in a common ancestor of 'S. carolinensis' and other closely related species.
The mutation that caused black fur happened independently in 'S. carolinensis' and all other closely related species.
The phenotypic variation shown in 'S. carolinensis' and other closely related species is caused by genetic drift.



The mutation that caused the C^B allele was due to a 24 base-pair deletion from the C^G allele.

0 1.4

The protein coded for by the C^B allele is 306 amino acids long.

Calculate the percentage reduction in size of the protein coded for by the C^B allele compared with the protein coded for by the C^G allele.

Give your answer to 3 significant figures and show your working. [2 marks]



Answer ____



In 'S. carolinensis', fur colour depends on the distribution and relative amounts of light pigments and dark pigments in the hairs of the fur. FIGURE 1, on page 16, shows how the protein produced from the C^G allele can result in the production of a light pigment or a dark pigment.



BLANK PAGE



FIGURE 1

C^G allele codes for the production of a receptor protein specific to a hormone called αMSH

αMSH binds to glutamic acid in the receptor protein

A protein called ASIP binds competitively to the receptor protein

αMSH leaves the receptor protein αMSH is blocked from binding to the receptor protein

The receptor protein becomes activated

The receptor protein is not activated

A dark pigment is produced

A light pigment is produced



The deletion mutation in the C^B allele results in the production of a receptor protein that does not have glutamic acid. The lack of glutamic acid in the receptor protein has the same effect as αMSH leaving the receptor protein.

|--|

Use FIGURE 1 and this information to suggest why 'S. carolinensis' with the genotype C^BC^B have black fur rather than grey fur. [3 marks]



		9



BLANK PAGE



0 2 . 1

Describe how the human immunodeficiency virus (HIV) is replicated ONCE INSIDE helper T cells (T _H cells). [4 marks]	





HIV-1 is the most common type of HIV. HIV-1 binds to a receptor on T_H cells called CCR5.

Current treatment for HIV-1 involves the use of daily antiretroviral therapy (ART) to stop the virus being replicated. Only 59% of HIV-positive individuals have access to ART.

Scientists have found that two HIV-1-positive patients (P and Q) have gone into remission (have no detectable HIV-1). This happened after a blood stem cell transplant (BSCT).

- Patient P was given TWO BSCTs, and patient Q was given ONE BSCT.
- All BSCTs came from a donor with T_H cells WITHOUT the CCR5 receptor.
- In addition, patient P had radiotherapy, and patient Q had chemotherapy. Both of these treatments are toxic.



 BOTH patients (P and Q) stopped receiving ART 16 months after BSCT.

18 months after stopping ART, BOTH patients had NO HIV-1 RNA in their plasma, NO HIV-1 DNA in their T_H cells and NO CCR5 on their T_H cells.

0 2 . 2

Use the information given to evaluate the use of BSCT to treat HIV infections.
[5 marks]





0 3


Scientists investigated movement in adult pine beetles. Adult beetles emerge from cracks in tree bark.

The scientists released a newly emerged adult beetle, G, from the centre of a sample area that had a single light source coming from one direction. They made a drawing of the beetle's path of walking. They repeated this with three more beetles, J, P and R.

FIGURE 2, on page 26, shows the scientists' results.



FIGURE 2



KEY



 Point from which beetle was released



0	3	1
		•

Name the type of behaviour shown by beetles G, J, P and R, and suggest ONE advantage to adult beetles of the type of behaviour shown. [2 marks]

Behaviour __		
Advantage		



At higher temperatures and higher light intensities, adult pine beetles normally


- move more
- fly rather than walk.

When preparing to fly, these adult beetles walk slowly. The scientists investigated the movement of adult beetles at different temperatures, and in the light and the dark. They created a box that was half in the light and half in the dark. They released an adult beetle at the midpoint of the central dividing line between light and dark areas. They recorded the path of the beetle's movement and its location after 5 minutes. From this, they calculated the mean speed of movement. They repeated the experiment with many beetles and at several temperatures.

FIGURE 3, on the opposite page, shows the scientists' results.



FIGURE 3





BLANK PAGE



0 3 . 2

After studying these experiments, a student concluded:

- there is a significant change in movement between 35 °C and 37.5 °C
- between 35 °C and 37.5 °C, more beetles move away from the light
- between 35 °C and 37.5 °C, more beetles have a slower walking speed.

might NOT be valid. [3 marks]							



BLANK PAGE



0 4

Freshwater marshes have one of the highest rates of gross primary production (*GPP*) and net primary production (*NPP*) of all ecosystems.

Carbon use efficiency (CUE) is the ratio of NPP:GPP. Freshwater marshes have a high CUE.



0	4	•	1
---	---	---	---

Use your knowledge of *NPP* to explain why freshwater marshes have a high *CUE* AND the advantage of this.

Do NOT refer to abiotic factors in your answer. [2 marks]

Explanation		
Auvantage_		



0 4		2
-----	--	---

Freshwater marsh soils are normally waterlogged. This creates anaerobic conditions.

Use your knowledge of the nitrogen cycle to suggest why these soils contain relatively high concentrations of ammonium compounds and low concentrations of nitrite ions and nitrate ions. [2 marks]



	 -	



A student investigated the growth rate of a freshwater marsh plant.

The growth rate (R) of a plant can be determined using this equation.

$$R = \frac{(\ln W_2 - \ln W_1)}{t}$$

Where

In = natural logarithm

t = duration of the investigation in days

 W_1 = plant biomass at the start of the investigation

 W_2 = plant biomass at the end of the investigation

The student used the equation above; however, she substituted height for biomass. This was because she did not want to destroy the plants to measure their biomass.



U 4 . 3	0	4		3
---------------	---	---	--	---

State the assumption the student has made AND suggest why this assumption might NOT be valid. [2 marks]				



04.4

At the end of the investigation, the student noted the freshwater marsh plant had grown 268 mm in height, and now measured 387 mm. She calculated the rate of growth (R) to be 0.097 mm m⁻¹ day⁻¹

Use this information and, SUBSTITUTING HEIGHT FOR BIOMASS, use the equation to calculate the duration of the student's investigation.

Give your answer to the nearest full day. Show your working. [2 marks]



____ days

[Turn over]

8



0	5		1
		_	_

The action of endopeptidases and exopeptidases can increase the rate of protein digestion. Describe how.
[2 marks]



n	5	2
U	J	

As humans age, there is a decrease in body protein.

Give the name of ONE body protein that could have resulted in:

[2 marks]

reduced muscle power	
reduced immunity	



Scientists investigated the effect of two types of dietary protein on the ability of old men to produce body proteins.

TABLE 2, on the opposite page, shows information about the two types of dietary protein investigated.



TABLE 2


Physiological factor	Name of dietary protein		
Tactor	Casein	Whey	
Rate of absorption of dietary protein / mmol dm ⁻³ amino acids in blood plasma h ⁻¹	3.05	4.33	
Stimulation of protein synthesis	Higher rate	Lower rate	
Breakdown of body proteins	No effect	Inhibitory effect	



FIGURE 4 shows the percentage of protein absorbed that becomes body protein in old men following a meal of casein or whey.

FIGURE 4

Percentage of protein absorbed that becomes body protein



A statistical test confirmed that the difference between the results shown in FIGURE 4 was significant.



0	5	3

would be cause a l	which type better for NET gain of mation properties.	r old men of body p ovided to	to eat to	Use





BLANK PAGE



0 6

Plants transport sucrose from leaves to other tissues for growth and storage. SUT1 is a sucrose co-transporter protein.

Scientists investigated whether the cells of tobacco plant leaves used SUT1 to transport sucrose to other tissues.



0	6		1
---	---	--	---

The scientists used a radioactively labelled DNA probe to show that the cells of tobacco plant leaves contained the *SUT1* gene.

Describe how they would do this.

Do NOT include PCR in your answer.
[4 marks]



	_



BLANK PAGE



06.2

To study the role of SUT1 in tobacco plants, scientists reduced the expression of the *SUT1* gene.

When the *SUT1* gene is transcribed, the SUT1 mRNA produced is called 'sense' SUT1 mRNA. The scientists genetically modified plants by inserting an EXTRA gene so that this ALSO allowed the production of 'antisense' SUT1 mRNA.

The scientists had two types of tobacco plants:

- type A plants that were genetically modified
- type B plants that were NOT genetically modified.



'antise would	Suggest now the production of fantisense' SUT1 mRNA in type A plants would reduce the expression of the SUT1 gene. [4 marks]					
						_



	_



06.3

The scientists hypothesised that lower rates of sucrose transport from leaves would cause reduced growth.


To test this hypothesis, the scientists provided leaves of type A and type B plants with labelled carbon dioxide (¹⁴CO₂). To estimate sucrose transport out of leaves, they measured the percentage of ¹⁴C remaining in the leaves for 16 hours.

FIGURE 5, on page 58, shows their results.



FIGURE 5

Percentage of ¹⁴C remaining in the leaves






Calculate the ratio of percentage of ¹⁴C remaining in leaves of type B to type A plants 16 hours after providing ¹⁴CO₂ [1 mark]



REPEAT OF FIGURE 5

Percentage of ¹⁴C remaining in the leaves





		4
U	b	4

In type B plants, the percentage of ¹⁴C remaining in the leaves does not reach zero per cent, as shown in FIGURE 5.

Suggest TWO reasons why. [2 marks]					
1					
2					



The scientists measured physiological differences between type A plants and type B plants.

TABLE 3 shows the scientists' results as they presented them.

TABLE 3

Physiological factor	Type of tobacco plant		
	Type A	Type B	
Rate of sucrose			
transport from leaf	0.1	3.7	
cells / µmol m ⁻² s ⁻¹			
Leaf sucrose			
concentration /	22	4	
mmol m ⁻²			
Ratio of shoot:root	6:1	2:1	
dry mass	0.1	Z. I	
Rate of			
photosynthesis / µmol	4	14	
glucose m ⁻² s ⁻¹			



Sucrose is able to inhibit the production and activity of rubisco in leaves of a plant. Type A plants have decreased dry mass compared with type B plants.

0	6	•	5
---	---	---	---

Use all the information to suggest AND explain how the physiological factors in TABLE 3 would contribute to the decreased dry mass observed in type A plants. [4 marks]





[Turn over]	15



SECTION B

Answer ONE question.

You are advised to spend no more than 45 minutes on this section.

0 7

Write an essay on ONE of the topics below.

EITHER

07.1

The importance of complementary shapes of molecules in organisms [25 marks]



OR

0	7	•	2
---	---	---	---

The importa	ance of ions	in metabolic
processes	[25 marks]	

















	_





















	_

















END OF QUESTIONS



Additional page, if required. Write the question numbers in the left-hand margin.



Additional page, if required. Write the question numbers in the left-hand margin.



BLANK PAGE

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
TOTAL		

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2021 AQA and its licensors. All rights reserved.

IB/M/CD/Jun21/7402/3/E2



