

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

AS

FURTHER MATHEMATICS

Paper 2 Mechanics

Time allowed: 1 hour 30 minutes

Materials

- You must have the AQA Formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Statistics). You will have 1 hour 30 minutes to complete **both** papers.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 40.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



Answer **all** questions in the spaces provided.

- 1** A light spring of natural length 0.6 metres is compressed to a length of 0.4 metres by a force of 20 newtons.

The stiffness of the spring is $k \text{ N m}^{-1}$

Find k

Circle your answer.

[1 mark]

20

50

100

200

- 2** State the dimensions of force.

Circle your answer.

[1 mark]

MLT

ML^2T

MLT^{-1}

MLT^{-2}



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



6 A ball of mass 0.15 kg is hit directly by a vertical cricket bat.

Immediately before the impact, the ball is travelling horizontally with speed 28 m s^{-1}

Immediately after the impact, the ball is travelling horizontally with speed 14 m s^{-1} in the opposite direction.

6 (a) Find the magnitude of the impulse exerted by the bat on the ball.

[2 marks]

6 (b) In a simple model the force, F newtons, exerted by the bat on the ball, t seconds after the initial impact, is given by

$$F = 10kt(0.05 - t)$$

where k is a constant.

Given the ball is in contact with the bat for 0.05 seconds, find the value of k

[3 marks]

Turn over ►



7 Use g as 9.81 m s^{-2} in this question.

A light elastic string has one end attached to a fixed point A on a smooth plane inclined at 25° to the horizontal.

The other end of the string is attached to a wooden block of mass 2.5 kg , which rests on the plane.

The elastic string has natural length 3 metres and modulus of elasticity 125 newtons .

The block is pulled down the line of greatest slope of the plane to a point 4.5 metres from A and then released.

7 (a) Find the elastic potential energy of the string at the point when the block is released.

[1 mark]

7 (b) Calculate the speed of the block when the string becomes slack.

[4 marks]



7 (c)

Determine whether the block reaches the point *A* in the subsequent motion, commenting on any assumptions that you make.

[3 marks]

Turn over for the next question

Turn over ►

8 Two spheres A and B are free to move on a smooth horizontal surface.

The masses of A and B are 2 kg and 3 kg respectively.

Both A and B are initially at rest.

Sphere A is set in motion directly towards sphere B with speed 4 m s^{-1} and subsequently collides with sphere B

The coefficient of restitution between the spheres is e

8 (a) (i) Show that the speed of B immediately after the collision is

$$\frac{8(1+e)}{5}$$

[4 marks]

8 (a) (ii) Find an expression, in terms of e , for the velocity of A immediately after the collision.

[2 marks]



8 (b) It is given that the spheres both move in the **same** direction after the collision.

Find the range of possible values of e

[2 marks]

8 (c) (i) The impulse of sphere A on sphere B is I
The impulse of sphere B on sphere A is J

Given that the collision is perfectly inelastic, find the value of $I + J$

[1 mark]

8 (c) (ii) State, giving a reason for your answer, whether the value found in part **(c)(i)** would change if the collision was **not** perfectly inelastic.

[2 marks]

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2021 AQA and its licensors. All rights reserved.



1 6

Jun21/7366/2M



2 1 6 A 7 3 6 6 / 2 M