



Cambridge International AS & A Level

CANDIDATE
NAME


--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/23

Paper 2 Pure Mathematics 2

October/November 2023

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.

- 3 (a) Find $\int_4^{10} \frac{4}{2x-5} dx$, giving your answer in the form $\ln a$, where a is an integer. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the exact value of $\int_4^{10} e^{2x-5} dx$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 4 (a) Sketch, on the same diagram, the graphs of $y = |3x - 5|$ and $y = 2x + 7$. [2]

- (b) Solve the equation $|3x - 5| = 2x + 7$. [3]

.....

.....

.....

.....

.....

.....

- (c) Hence solve the equation $|3^{y+1} - 5| = 2 \times 3^y + 7$, giving your answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

- 6 (a) Show that $\operatorname{cosec} \theta(3 \sin 2\theta + 4 \sin^3 \theta) \equiv 4 + 6 \cos \theta - 4 \cos^2 \theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Solve the equation

$$\operatorname{cosec} \theta(3 \sin 2\theta + 4 \sin^3 \theta) + 3 = 0$$

for $-\pi < \theta < 0$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find $\int \operatorname{cosec} \theta (3 \sin 2\theta + 4 \sin^3 \theta) \, d\theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Show that $q = \sqrt[3]{2 + 18 \ln 3 - q}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show by calculation that the value of q lies between 2.5 and 3.0. [2]

.....

.....

.....

.....

.....

- (d) Use an iterative formula, based on the equation in (b), to find the value of q correct to 4 significant figures. Give the result of each iteration to 6 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

