Surname

Centre Number

2

Other Names

wjec

GCE AS/A LEVEL – NEW

2400U20-1

BIOLOGY – AS unit 2 Biodiversity and Physiology of Body Systems

TUESDAY, 6 JUNE 2017 – AFTERNOON

1 hour 30 minutes

For Examiner's use only			
Question	Maximum Mark	Mark Awarded	
1.	10		
2.	11		
3.	10		
4.	13		
5.	15		
6.	12		
7.	9		
Total	80		

ADDITIONAL MATERIALS

In addition to this examination paper, you will need a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer all questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation pages at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The assessment of the quality of extended repsonse (QER) will take place in question **7**. The quality of written communication will affect the awarding of marks.




BLANK PAGE

2


PLEASE DO NOT WRITE ON THIS PAGE







(b) The diagram below shows phylogenetic trees illustrating the coevolution of different species of lice and their vertebrate hosts. The dotted lines represent parasite-host relationships. The numbers at branch nodes on each tree represent the estimated time of divergence from a common ancestor in millions of years ago (MYA). These times were obtained using a molecular clock supported by fossil evidence. A molecular clock measures the degree of genetic similarity between species.





© WJEC CBAC Ltd.

Examiner only

Examiner only

- Suggest how DNA analysis could be used to measure the degree of genetic similarity between species.
- (iii) The table below represents a hypothetical molecular clock. Insert the value for the time of divergence for 90% genetic similarity in the table below. [1]

Degree of genetic similarity / %	Time of divergence from a common ancestor / MYA
40	30
60	20
80	10
90	


- (c) Another human ectoparasite is the disease-carrying body louse. The body louse attaches its eggs to clothing whereas the head louse attaches its eggs to scalp hair. Suggest why they are considered to be the same species.

10

2400U201 05










Examiner only The photomicrograph below is a cross section through the trachea and oesophagus of a (b) mammal. oesophagus trachea The rings of chitin in an insect's trachea are complete. The rings of cartilage in a mammal's trachea are incomplete or "C-shaped". Use the information in the (i) photomicrograph to suggest why the rings of cartilage in a mammal's trachea are incomplete. [1] Name one plant tissue that shows a similar pattern of support material to that seen (ii) in an insect's tracheae. [1]

7



Turn over.

2400U201 07





	9	
(ii)	Suggest one change that you would expect to see in these curves during strenuous exercise. [1]	Examiner only
		11
		2400U201



2400U201 09

 and potato leafroll. Elm bark beetles are vectors of Dutch elm disease. These beetles are often contaminate with fungal spores gain entry to xylem vessels. The spores germinate and produce a mycelium which leads to the blockage of xylem vessels. The earliest external symptoms of infectio are chlorosis (yellowing) and wilting of leaves above the infection site. These leaves ofte turn brown and curl up. Symptoms often spread rapidly leading to the death of the tree. Potato leafroll is caused by a virus carried by aphids. When aphids feed, the virus enter the phloem. The virus infects cells in the leaves and the roots. Symptoms include chloros and rolling of leaves, and death of potato tuber cells. (i) Explain why blockage of xylem vessels by the fungus causes the wilting of leaver above the infection site and the death of elm trees. (ii) Explain the difference in the distribution of the symptoms of potato leafroll in a plate. 		
 with fungal spores. When the beetles feed on the young bark of healthy elm trees th fungal spores gain entry to xylem vessels. The spores germinate and produce a mycelium which leads to the blockage of xylem vessels. The earliest external symptoms of infectic are chlorosis (yellowing) and wilting of leaves above the infection site. These leaves ofte turn brown and curl up. Symptoms often spread rapidly leading to the death of the tree. Potato leafroll is caused by a virus carried by aphids. When aphids feed, the virus enter the phloem. The virus infects cells in the leaves and the roots. Symptoms include chloros and rolling of leaves, and death of potato tuber cells. (i) Explain why blockage of xylem vessels by the fungus causes the wilting of leave above the infection site and the death of elm trees. (ii) Explain the difference in the distribution of the symptoms of potato leafroll in a plain 	(a)	Some insects are vectors of plant diseases. Two such diseases are Dutch elm disease and potato leafroll.
 the phloem. The virus infects cells in the leaves and the roots. Symptoms include chloros and rolling of leaves, and death of potato tuber cells. (i) Explain why blockage of xylem vessels by the fungus causes the wilting of leave above the infection site and the death of elm trees. 		Elm bark beetles are vectors of Dutch elm disease. These beetles are often contaminated with fungal spores. When the beetles feed on the young bark of healthy elm trees the fungal spores gain entry to xylem vessels. The spores germinate and produce a mycelium which leads to the blockage of xylem vessels. The earliest external symptoms of infection are chlorosis (yellowing) and wilting of leaves above the infection site. These leaves ofter turn brown and curl up. Symptoms often spread rapidly leading to the death of the tree.
(ii) Explain the difference in the distribution of the symptoms of potato leafroll in a plai		Potato leafroll is caused by a virus carried by aphids. When aphids feed, the virus enters the phloem. The virus infects cells in the leaves and the roots. Symptoms include chlorosis and rolling of leaves, and death of potato tuber cells.




11 Examiner only The photomicrograph below is a transverse section of phloem tissue taken using a transmission electron microscope. (b) С A В 2400U201 11 Identify the cells labelled A and B. [1] (i) Α Β..... Give one reason for your choice of answer to (b)(i). (ii) [1]




Examiner only The photomicrograph below is of region ${\bf C}$ at a higher magnification. It shows the adjoining cell walls of ${\bf A}$ and ${\bf B}.$ (iii) Identify structures **D** and explain their importance in the functioning of phloem. [2] 10









© WJEC CBAC Ltd.


14

.td.

(i)	During the investigation the temperature was maintained constant at 20 °C. Name two other environmental factors that should have been kept constant during the investigation. [2]
(ii)	Which unit should have been included on the bar chart to represent the mean transpiration rate?
(iii)	The student concluded that the transpiration rates of the two plants were significantly different. Use the information in the bar chart to explain why she was confident in reaching this conclusion. [1]
(iv)	Suggest one structural adaptation of navelwort and explain how it could account for the difference in transpiration rates of privet and navelwort. [2]
(v)	The student found that the scale on the capillary tube was accurate to ±1.0 mm. She measured the time taken for the air bubble to travel a distance of 50 mm along the capillary tube. Calculate the percentage error of the equipment over this distance. Show your working in the space below. [2]



Turn over.



© WJEC CBAC Ltd.

The table shows the partial pressures of oxygen (pO2) and carbon dioxide (pCO2) at (b) different sites in the human body. Site pO₂ / kPa pCO₂ / kPa lungs 10 2 muscle tissue fluid at 4 4 rest muscle tissue fluid 1 8 during exercise The graph shows that at the pO_2 and pCO_2 typical in the lungs the percentage (i) saturation of haemoglobin with oxygen is 96%. Give the percentage saturation of haemoglobin with oxygen in muscle tissue fluid when it is: [2] at rest%% during exercise Explain the significance of the effect of an increase in pCO₂ for respiring muscle (ii) tissue. [2] Respiratory minute volume is the volume of gas inhaled or exhaled from a person's lungs (C) per minute. The minute volume of a healthy person during normal breathing at rest is $6 - 7 \text{ dm}^3 \text{min}^{-1}$. In people with chronic diseases, such as heart disease, the minute volume is $12 - 16 \text{ dm}^3 \text{min}^{-1}$. Explain how an increase in minute volume results in a decrease in pCO₂ of blood in (i) alveolar capillaries. [2] Suggest why people with reduced blood pCO₂ commonly feel tired and lack energy. (ii) [2]


(2400U20-1)



© WJEC CBAC Ltd.

Turn over.

Examiner only (*d*) Most carbon dioxide is carried as hydrogen carbonate ions (HCO₃⁻) in the plasma. The following chemical pathway shows how carbon dioxide is converted into HCO₃⁻ in a red blood cell.



(iii) The table shows the concentrations of hydrogen carbonate ions and chloride ions in the blood plasma of an arteriole entering and a venule leaving a respiring muscle.

Pland variable	Plasma concentration / mmol dm ⁻³	
Blood vessel	Hydrogen carbonate ions	Chloride ions
Arteriole	22	106
Venule	30	98

Explain the changes in the concentration of chloride ions as shown by the table above. [3]

18

Examiner

© WJEC CBAC Ltd.

BLANK PAGE

19

PLEASE DO NOT WRITE ON THIS PAGE



6. Pepsin and trypsin are enzymes involved in the digestion of proteins in the alimentary canals of mammals. Both enzymes are endopeptidases.

A group of students performed an experiment to investigate the effect of an endopeptidase, an exopeptidase and a mixture of the two enzymes on the digestion of protein in milk powder. As the protein is digested the solution becomes clear.

Equal volumes and concentrations of the enzyme solutions were added to equal volumes and concentrations of milk powder solution. The pH was maintained using a pH 8 buffer solution. The time taken for the solution to become clear was recorded, as shown below.

	Time taken for milk solution to become clear / s			
Student group	Endopeptidase	Exopeptidase	Mixture of endopeptidase and exopeptidase	
A	43	74	19	
В	77	95	39	
С	69	93	34	
D	88	138	34	
E	52	69	36	
F	47	71	34	
G	109	198	60	
Н	61	80	42	
I	83	166	26	
J	38	60	15	
Mean for all groups		104	34	

(a) Calculate the mean time for the milk solution to become clear when mixed with the endopeptidase. **Insert your answer in the table above**. [2]

(b) The results show a great deal of variation. However, it was decided that there were no anomalous results. Explain why this decision was made. [1]




(c)	Suggest two sources of inaccuracy within the experimental method, which could account for the variation in the results of the groups. For each of your suggestions give one way in which the method could be improved. [4]	Examiner only
······		
······		
(d)	Explain why digestion was more rapid when using the enzyme mixture than when using the individual enzymes. [3]	
······		
(e)	Explain why pepsin was not the endopeptidase chosen for use in this investigation. [2]	
		12



Turn over.

(2400U20-1)

Turr



Examiner only



Examiner only



Examiner only 9 END OF PAPER





PLEASE DO NOT WRITE ON THIS PAGE



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only
		1
	······	



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examine only

